Eurasia Specialized Veterinary Publication

Entomological Research Letters

ISSN: 3108-639X

2024, Volume 4, Issue 2, Page No: 93-103
Copyright CC BY-NC-SA 4.0
Available online at: www.esvpub.com/

Post-Transcriptional RNA Regulation Extends Molecular Control Beyond Gene Expression During Insect Development and Environmental Stress

Siti Rahman[1]*, Ahmad Zaki[1], Nurul Huda[2], Amir Faisal[1], Lim Wei[2]

[1]Department of Post-Transcriptional Regulation and Development, Faculty of Life Sciences, University of Malaya, Kuala Lumpur, Malaysia.
[2]Department of RNA Control and Environmental Stress, Faculty of Biosciences, Universiti Teknologi Malaysia, Johor Bahru, Malaysia.

***E-mail** ✉ siti.rahman@um.edu.my

ABSTRACT

Insect development and environmental responses are often interpreted through transcriptional change, yet transcript abundance represents only one stage of molecular regulation. Post-transcriptional processes can influence whether transcripts persist, which isoforms are produced, how RNA information is modified, which molecules are targeted by small RNAs, and how efficiently transcripts are translated. This conceptual analysis examines RNA stability and degradation, alternative splicing, RNA editing, small RNAs, RNA modifications, and translational regulation as distinct but potentially interacting regulatory dimensions. The article argues that post-transcriptional control should not be interpreted as a single pathway but as a set of conditional regulatory states whose contribution depends on developmental stage, taxonomic context, environmental exposure, and measurement strategy. A proposed integrative framework distinguishes directly demonstrated molecular mechanisms from broader synthesis and identifies boundaries where transcript-level observations cannot alone support organismal or evolutionary conclusions. By organizing evidence across regulatory layers, this article provides a conceptual basis for interpreting insect molecular responses beyond gene expression measurements while maintaining clear limits on causal inference.

Keywords: Insects, Post-transcriptional regulation, RNA stability, Alternative splicing, RNA editing, Epitranscriptomics

Received: 28 September 2024
Revised: 15 November 2024
Accepted: 17 November 2024

How to Cite This Article: Rahman S, Zaki A, Huda N, Faisal A, Wei L. Post-Transcriptional RNA Regulation Extends Molecular Control Beyond Gene Expression During Insect Development and Environmental Stress. Entomol Res Lett. 2024;4(2):93-103. https://doi.org/10.51847/cXpz0rldCq

Introduction

Gene expression studies have transformed insect biology by revealing transcriptional responses associated with development, environmental variation, and physiological transitions. However, RNA abundance is not equivalent to biological output. Between transcription and protein production, multiple regulatory processes determine transcript persistence, molecular identity, accessibility, and translation. The increasing availability of transcriptomic, ribosome-profiling, and epitranscriptomic approaches has therefore expanded the question from whether genes are expressed to how RNA molecules are processed after synthesis.

A central challenge is that post-transcriptional regulation is often discussed as a broad concept despite representing multiple mechanistically distinct processes. Small RNAs, alternative splicing, RNA modification, RNA editing, RNA decay, and translational control operate at different molecular stages and are measured through different experimental approaches. Evidence from insect systems demonstrates that these mechanisms can participate in developmental regulation, but the strength of inference differs substantially among profiling studies, functional perturbations, biochemical validation, and organism-level phenotyping. Functional interpretation of insect small-

© 2024 Entomological Research Letters

RNA systems, for example, increasingly depends on combining expression patterns with target validation and experimental manipulation rather than relying on sequence-based predictions alone [1].

Insect development provides particularly informative examples because hormonal transitions, tissue remodeling, and environmental sensitivity require precise temporal regulation. Reviews of insect post-transcriptional processes have shown that microRNAs, alternative splicing, and endocrine pathways can interact during metamorphosis and reproduction, but the degree of integration differs among taxa and developmental contexts [2]. Developmental plasticity studies further demonstrate that transcript abundance, differential splicing, and microRNA-associated patterns may change simultaneously while representing distinct regulatory dimensions rather than a single coordinated mechanism [3].

The relationship between RNA abundance and downstream molecular output also requires caution. Time-resolved transcriptomic and proteomic analysis in Drosophila demonstrated that protein trajectories cannot be interpreted simply from RNA measurements because translation and degradation dynamics contribute substantially to observed patterns [4]. Therefore, the central problem addressed here is not whether post-transcriptional regulation exists, but how molecular control should be interpreted when multiple RNA regulatory layers intervene between transcription and phenotype.

This article develops a conceptual analysis of six post-transcriptional domains: RNA stability and degradation, alternative splicing, RNA editing, small RNAs, RNA modifications, and translational regulation. The proposed synthesis does not assume that these mechanisms form a universal pathway. Instead, it organizes evidence around regulatory states, mechanistic boundaries, and inference limits.

Results and Discussion

RNA stability represents one of the earliest post-transcriptional decision points after transcript production. A transcript may persist, decay rapidly, or change stability according to sequence features, RNA-binding proteins, cellular state, and developmental context. Studies in Drosophila demonstrate that RNA persistence can be actively regulated rather than treated as a passive consequence of transcription.

Orb2 provides one example where transcript stability is linked to developmental state. During neuronal differentiation, Orb2-associated regulation influences expression of rare-codon-enriched transcripts, demonstrating that sequence composition can participate in context-dependent RNA stabilization [5]. This finding does not establish a universal relationship between codon usage and RNA stability across insects, but it illustrates that transcript persistence can depend on regulatory machinery operating after transcription.

Direct biochemical evidence further demonstrates that RNA decay can be mechanistically controlled. Smaug and Cup recruit CCR4-NOT-dependent deadenylation machinery to nanos messenger RNA in a reconstituted Drosophila system, showing a defined molecular pathway linking RNA-binding proteins with transcript degradation [6]. Because this evidence derives from a specific maternal RNA regulatory system, it provides a mechanistic example rather than evidence for a universal insect decay program.

RNA modifications can also influence stability. In early Drosophila embryos, m^6A modification contributes to FMR1-associated RNA granule behavior and maternal RNA decay, linking chemical RNA state with transcript fate [7]. This connection demonstrates that RNA stability can depend on molecular context beyond transcript sequence alone, although the identified mechanism applies to defined targets rather than all modified transcripts.

Genome-wide analyses further indicate that RNA stability has conditional effects. Stability changes may buffer transcriptional variation or reinforce differential expression depending on biological context [8]. Therefore, RNA stability should not be interpreted simply as an amplification mechanism. It may either stabilize molecular states or permit rapid regulatory transitions.

Alternative splicing

Alternative splicing expands transcript diversity by allowing a single genomic locus to generate multiple RNA isoforms. In insects, this mechanism contributes to developmental regulation, environmental responses, and tissue-specific molecular states. However, the presence of differential splicing does not automatically indicate functional adaptation or improved performance.

Seasonal plasticity studies in butterflies demonstrate that differential splicing can represent a regulatory response distinct from differential gene expression. In Bicyclus anynana, environmentally associated developmental variation involved both transcript abundance changes and alternative splicing patterns, indicating that isoform

regulation provides information not captured by expression measurements alone [9]. The evolutionary consequences of these changes remain context dependent because molecular plasticity does not by itself demonstrate adaptive value.

Developmental timing also influences splicing patterns. Honeybee embryonic analysis revealed that sex- and stage-dependent splicing differences contribute to developmental molecular variation within a haplodiploid system [10]. These findings highlight the importance of life-history context because splicing patterns observed during one developmental interval cannot automatically be transferred to other insect stages.

Functional studies provide stronger evidence when specific isoforms are experimentally manipulated. In Bombyx mori, alternative splicing of POUM2 contributes to embryonic cuticle formation and tanning, demonstrating that isoform identity can affect developmental outcomes in a defined system [11]. Similarly, long-read sequencing approaches reveal that alternative splicing can interact with alternative polyadenylation, showing that RNA processing events may be connected within individual transcripts rather than acting as completely independent layers [12].

Alternative splicing may also contribute to developmental robustness. Manipulation of Yorkie isoform balance in Drosophila demonstrates that specific splicing states influence developmental stability [13]. However, developmental stability should not be interpreted as evidence of evolutionary adaptation without direct fitness measurements.

RNA editing

RNA editing introduces sequence changes after transcription and therefore creates a regulatory layer in which RNA information can differ from genomic information. In insects, editing patterns vary across taxa, developmental states, tissues, and environmental conditions, making interpretation highly dependent on experimental context.

The first A-to-I editome analysis in the hemipteran Coridius chinensis demonstrated extensive editing variation and revealed that cold exposure could reduce global editing while increasing editing at selected sites [14]. This opposing pattern illustrates that environmental regulation of RNA editing is not necessarily directional. A reduction in total editing does not imply reduced regulatory importance because individual sites may respond differently.

Interpretation of editing landscapes also depends strongly on detection methods. Slic-seq provides a methodological approach for transcriptome-wide A-to-I editing profiling and illustrates the importance of validation strategies when distinguishing true editing events from technical artifacts [15].

Natural variation further demonstrates that RNA editing can reflect interactions between genetic background and environment. In Drosophila populations from divergent microclimates, editing variation was associated with both genetic and temperature-related factors, including a validated cis-regulatory example [16]. However, environmental association does not alone establish that editing differences are adaptive.

ADAR biology also requires separation of editing-dependent and editing-independent effects. Genetic studies show that ADAR contributes to neural and immune functions through both catalytic editing activity and additional protein-mediated roles [17]. Therefore, phenotypes associated with ADAR manipulation cannot automatically be attributed exclusively to RNA editing.

The interpretation of RNA editing requires distinguishing at least four analytical states: observed editing change, mechanistic editing consequence, organism-level phenotype, and evolutionary interpretation. These states should not be merged without direct evidence.

The different evidence patterns surrounding RNA editing require an analytical map rather than a simple list of mechanisms. The table below is designed to distinguish molecular observation from interpretation and from unresolved inference boundaries.

Table 1. Analytical states determining how insect RNA editing evidence can be interpreted

Editing state	Biological provenance	Observable molecular basis	Consequence that can be supported	Analytical ceiling	Required validation
Environmental and site-specific signal					
Global editing shift under environmental exposure	Defined insect stress systems	Transcriptome-wide editing comparison	Indicates altered editing landscape	Direction may differ among sites	Validate candidate sites functionally

Site-specific recoding response	Individual genes or transcripts	Matched DNA/RNA editing analysis	May indicate regulated transcript alteration	Functional consequence may remain unknown	Prioritize experimentally tested sites
Genotype and discovery-method dependence					
Genotype-associated editing variation	Natural populations	Population genomic and transcriptomic comparisons	Indicates interaction between genetic background and RNA regulation	Does not establish selection	Combine ecological and functional approaches
Method-dependent editing discovery	Editing detection workflows	Validation-dependent computational and experimental pipelines	Determines confidence in detected events	Technical assumptions influence calls	Apply orthogonal confirmation
Cross-layer synthesis					
Proposed cross-layer interpretation	Integrated RNA regulation concept	Synthesis across mechanisms	Suggests possible links between editing and broader molecular states	Not a universal insect mechanism	Test within shared experimental systems

Small RNAs

Small RNAs provide a regulatory mechanism through which short RNA molecules influence transcript stability, translation, and developmental timing. In insects, small-RNA effects are particularly important during developmental transitions and stress exposure, although the strength of evidence differs among systems.

Early environmental stress can influence maternal small-RNA states before later developmental processes become established. Heat-shock experiments in Drosophila embryos demonstrated stress-sensitive changes in maternal microRNAs and associated molecular consequences involving Dicer-1-dependent regulation [18]. However, downstream chromatin relationships require careful interpretation because molecular association does not automatically establish a complete causal chain.

Developmental timing provides stronger functional examples. In Bombyx mori, disruption of let-7 alters hormonal regulation of metamorphosis, demonstrating a causal role for a specific microRNA in developmental transition [19]. Similarly, miR-14 manipulation affects larval timing and ecdysteroid-associated processes, although individual target relationships require appropriate validation [20].

The assignment of functional microRNA targets requires more than sequence complementarity. Integrated AGO1 binding, expression, and translation analyses demonstrate that functional target identification benefits from combining multiple evidence types [21].

The biological meaning of small-RNA regulation therefore depends on the relationship between molecular detection, target confirmation, developmental context, and phenotype. To represent this reasoning visually without reducing it to a simple pathway, the following figure separates observed states, regulatory transitions, and inference boundaries.




Figure 1. Environmental and developmental states shaping insect small-RNA regulatory interpretation

RNA modifications

RNA modifications create an additional regulatory dimension by altering RNA properties without changing genomic sequence. In insects, epitranscriptomic regulation has been linked to development, environmental responses, endocrine regulation, and transcript processing. However, modification presence alone does not establish biological function.

In Plutella xylostella, m^{6}A regulation influences juvenile hormone-related processes under a defined resistance-associated challenge context. METTL3/14-dependent modification affects JHE expression and hormonal state, illustrating how RNA modification can connect molecular regulation with physiology [22]. This example demonstrates a specific context-dependent mechanism rather than a general rule that m^{6}A improves stress performance.

Developmental evidence also exists for other RNA modifications. Comparative analysis of maternal mRNA m^{5}C landscapes showed broad modification patterns across animals, while functional experiments in Drosophila demonstrated that loss of maternal m^{5}C affects developmental timing during early embryogenesis [23]. The comparative distribution of modification does not imply identical functional roles across insect species.

The conservation of RNA modification machinery provides another important distinction. Analysis across insect species revealed conservation of m^{6}A-related genes, while functional manipulation of METTL3 in Bombyx mori altered embryonic transcriptomic states [24]. These results support conservation of regulatory components but do not demonstrate identical downstream mechanisms among insects.

RNA modification can also intersect with other RNA-processing layers. In Drosophila, Hakai contributes to the m^{6}A modification pathway and influences Sxl-associated alternative splicing [25]. This provides direct evidence that some post-transcriptional mechanisms can interact, while leaving open how frequently such interactions occur across insect diversity.

The interpretation of RNA modifications requires distinguishing modification detection, molecular mechanism, organismal consequence, and broader evolutionary meaning. The following table addresses this distinction.

Table 2. Comparative interpretation of insect RNA modification states across molecular evidence levels

Physiological and maternal states		
RNA modification comparison dimension	**Modification linked to endocrine physiology**	**Maternal developmental modification state**
Biological context and evidence	Defined insect challenge systems Modification mapping plus hormonal assays	Early embryonic systems

		Epitranscriptomic mapping and functional perturbation
Inference enabled	Connects RNA state with physiological regulation	Supports developmental regulation
Unresolved boundary	Context-specific mechanism	Functional evidence concentrated in model insects
Decisive experiment	Test additional developmental and environmental settings	Expand comparative validation
Machinery and modification-splicing support		
RNA modification comparison dimension	**Conserved modification machinery**	**Modification–splicing interaction**
Biological context and evidence	Broad insect comparisons Gene-family and evolutionary analyses	Defined developmental pathways Modification assays plus splicing analysis
Inference enabled	Indicates molecular conservation	Demonstrates cross-layer regulation
Unresolved boundary	Conservation does not equal conserved function	Limited to studied systems
Decisive experiment	Compare downstream targets	Identify additional coupled processes
Integrated modification role		
RNA modification comparison dimension	**Proposed integrated modification role**	
Biological context and evidence	Conceptual synthesis Cross-study interpretation	
Inference enabled	Suggests modification as a regulatory state variable	
Unresolved boundary	Not an experimentally validated universal model	
Decisive experiment	Test shared systems experimentally	

Translational regulation

Transcript abundance does not determine protein output unless translation is also considered. Translational regulation therefore represents a terminal but independently regulated stage of post-transcriptional control. Its importance is especially clear when transcript architecture or ribosome state changes the probability that an available messenger RNA will generate protein.

In Drosophila, the RNA-binding protein Nocte promotes translation reinitiation on transcripts containing long upstream open reading frames. Genetic disruption, ribosome profiling, and rescue experiments link this activity to normal eye development, demonstrating that transcript availability can be separated from translational competence [26]. The mechanism is target restricted: not every Nocte-responsive transcript shares the same upstream-open-reading-frame organization, so the study does not justify a universal reinitiation rule.

Translation can also be regulated through the ribosome itself. OTUD6-dependent deubiquitination of RPS7 on free 40S ribosomal subunits alters global translation and responds to defined alkylation and translational-stress conditions in Drosophila [27]. This establishes ribosome biochemical state as a regulatory variable, but laboratory stress responses should not be equated with field tolerance or adaptive environmental performance.

Measurement adds another boundary. Developmental Poly-Ribo-seq in Drosophila embryos distinguished productive translation from reproducible ribosome association that does not necessarily yield canonical protein output [28]. Consequently, ribosome occupancy, like transcript abundance, is an intermediate observation rather than an automatically interpretable endpoint. Translation must be evaluated using assays capable of distinguishing engagement, reading-frame behavior, and productive protein synthesis when those distinctions matter.

These studies extend post-transcriptional control beyond RNA processing. Transcript sequence architecture, RNA-binding proteins, ribosome modification state, and developmental timing can alter whether an RNA molecule becomes functional protein. The proposed synthesis therefore treats translational realization as a separate regulatory dimension rather than the inevitable consequence of transcript presence.

Integrated post-transcriptional control across development and stress

The six regulatory domains considered here act on different properties of an RNA molecule: persistence, isoform identity, nucleotide information, small-RNA targeting, chemical state, and translational realization. Their coexistence does not mean that they constitute one linear pathway. Developmental transcriptomic data in Bicyclus, for example, show differential expression, splicing, and microRNA variation in the same developmental system without demonstrating that those changes form a single causal chain [3]. Likewise, m6A-sensitive maternal RNA decay in Drosophila establishes one direct connection between chemical modification and RNA fate, but not a general rule connecting every epitranscriptomic mark to degradation [7].

The evidence supports parallel regulation and bounded cross-layer coupling. RNA stability can either buffer or reinforce abundance changes [8]. Alternative splicing may depend on connectivity to alternative 3-prime ends in selected transcripts [12]. ADAR-associated phenotypes can contain editing-dependent and editing-independent components [17]. Functional microRNA regulation becomes more persuasive when binding information is combined with expression and translational consequences [21]. The m6A writer pathway can also intersect directly with sex-specific splicing in Drosophila [25]. These examples justify representing post-transcriptional regulation as a network of conditionally connected states, while resisting the stronger claim that every layer interacts in every insect.

A second distinction is scale. Molecular responses do not automatically establish organismal benefit or evolutionary adaptation. Seasonal alternative splicing in Bicyclus may be relevant to environmental plasticity, but its adaptive value cannot be inferred from transcript patterns alone [9]. Cold-associated RNA editing in Coridius likewise includes global and site-specific responses without demonstrating that those changes improve fitness [14]. Translation changes under defined laboratory stress demonstrate regulatory sensitivity, not field-level ecological performance [27].

This article therefore proposes a post-transcriptional state architecture with three explicit boundaries. The first separates molecular detection from validated mechanism. The second separates mechanism from organismal consequence. The third separates phenotype from evolutionary or ecological interpretation. A relationship crosses a boundary only when the evidence directly supports the stronger inference. Cross-layer links are treated as evidence-supported only when demonstrated within a compatible experimental system; otherwise they remain proposed synthesis.

Figure 2 reconciles mechanism-specific evidence across molecular and inferential levels, while **Table 3** records how post-transcriptional states alter interpretation without unsupported numerical scores.




Figure 2. Regulatory-state reconciliation across RNA processing, molecular output, and higher-order inference

Table 3. State-and-response ledger for interpreting integrated post-transcriptional control in insects

Regulatory layer	Biological setting	Molecular observation	Interpretive consequence	Boundary, cross-layer test, and state
Mechanism-specific control layers				
Persistence-dominant state	Developmental or perturbation context in which RNA stability changes independently of transcription	Transcript decay or stability measurement	Transcript abundance may be buffered or reinforced after synthesis	Stability effect is context dependent Pair transcriptional measurements with decay estimates Evidence-supported class
Isoform-selection state	Stage-, sex-, or tissue-dependent RNA processing	Full-transcript or splice-junction evidence	Functional RNA identity may change without equivalent total-expression change	Isoform function cannot be assumed from detection Validate isoform-specific activity Evidence-supported class
Editing-divergence state	Genetic or temperature-sensitive editing system	Matched DNA/RNA and site validation	RNA sequence state may differ from genomic sequence	Site response may not predict organismal benefit Test site-specific function and genetic dependence Evidence-supported class
Targeted small-RNA state	Developmental or stress-sensitive small-RNA system	Small-RNA abundance plus target evidence	Selected transcripts may change stability or translation	Prediction or binding alone is insufficient Combine perturbation with target-output assays Evidence-supported class
Modification-coupled state	Embryonic, endocrine, or transcript-processing context	Modification mapping plus functional perturbation	Chemical RNA state may alter decay, splicing, or physiology	Effects remain mark-, target-, and taxon-specific Test the same modification across matched contexts Evidence-supported class
Translation-gated state	Development or defined cellular stress	Ribosome engagement and productive-translation evidence	Protein output can diverge from transcript availability	Occupancy need not equal protein synthesis Resolve reading frame and productive output Evidence-supported class
Cross-layer regulatory state				
Cross-layer regulatory state	Same biological system contains a demonstrated interaction between two RNA layers	Mechanistic evidence connecting layers	One post-transcriptional process conditions another	Demonstrated examples do not establish universal coupling Replicate cross-layer mechanisms across taxa Evidence-supported only where directly demonstrated
Higher-order integration				
Higher-order integrated state	Multiple post-transcriptional signals occur in the same developmental or stress context	Cross-study or multi-omic synthesis	Multiple RNA states may jointly constrain molecular output	Causal topology and scale transfer remain unresolved Test competing network structures experimentally Proposed synthesis

Transcriptional measurements remain valuable but occupy one level of a larger regulatory system. Transcriptomics remains indispensable for identifying molecular responses. The problem arises when transcript abundance is treated as equivalent to protein output or biological mechanism. Time-resolved Drosophila analysis demonstrates that translation and degradation kinetics can account for substantial protein dynamics that would otherwise appear as unexplained transcript-protein discordance [4]. RNA-stability studies similarly show that post-transcriptional regulation may either oppose or reinforce transcriptional changes [8].

A useful interpretation of insect molecular regulation must therefore preserve the property being measured. Abundance, stability, isoform composition, nucleotide editing, small-RNA occupancy, RNA modification, ribosome engagement, and protein production are related observations, but they are not interchangeable. The proposed state architecture formalizes this distinction by asking which molecular property changed, what evidence establishes the mechanism, and how far the resulting inference can legitimately extend.

Environmental responsiveness requires especially careful language. Alternative splicing associated with seasonal plasticity may generate functionally relevant variation, yet molecular plasticity alone does not demonstrate adaptation [9]. Editing differences across divergent Drosophila microclimates reflect both environmental and genetic components, showing why observed population divergence cannot automatically be assigned to selection [16]. Stress-sensitive maternal microRNAs provide another example in which a defined embryonic perturbation alters molecular state, but the complete causal route to later chromatin effects remains incompletely resolved [18]. Similarly, the m6A–juvenile-hormone mechanism demonstrated in Bt-resistant Plutella occurs in a specific resistance-associated physiological context and should not be generalized to all environmental stress [22].

Taxonomic concentration is another major limitation. Mechanistic evidence is dominated by Drosophila, with important but fewer examples from Bombyx, Bicyclus, Plutella, Apis, and Coridius. The comparative review literature itself emphasizes the uneven functional resolution of small-RNA biology outside model insects [1]. The 207-species survey of m6A-related genes broadens phylogenetic coverage, but conservation of machinery does not establish conservation of downstream regulatory function [24]. These asymmetries make "insect post-transcriptional regulation" a useful comparative research domain, not a single conserved mechanism.

Future work should prioritize designs in which multiple regulatory layers are measured in the same specimens, developmental interval, or environmental manipulation. Long-read approaches can resolve transcript connectivity that short-read analyses may fragment [12]. Editing studies require orthogonal strategies that discriminate genuine nucleotide conversion from sequencing or mapping artifacts [15]. MicroRNA studies gain mechanistic strength when binding, perturbation, transcript response, and translational response converge [21]. Ribosome-profiling approaches likewise require interpretation of productive versus nonproductive engagement [28].

The strongest tests of the proposed architecture should compare competing explanations rather than accumulate molecular layers. A developmental phenotype attributed to post-transcriptional integration should be tested against simpler explanations based on transcription, RNA turnover, or one dominant processing event. Likewise, a stress-associated molecular state should be separated experimentally from generalized damage, acclimatory plasticity, developmental delay, and genetically structured population differences. Such tests would determine when post-transcriptional integration provides explanatory value beyond parallel molecular responses.

Experimental procedures: conceptual evidence selection and network construction

This article used a conceptual evidence-selection strategy rather than an empirical experiment, systematic review, or meta-analysis. Evidence was organized by regulatory mechanism and inferential strength. Studies were used to support claims about RNA stability, splicing, editing, small RNAs, RNA modifications, translation, methodological limitations, or cross-layer interactions only when their design directly addressed that claim.

The evidence hierarchy was qualitative rather than numerical. Profiling established molecular occurrence or association; matched molecular measurements strengthened state assignment; genetic perturbation supported functional inference; biochemical reconstitution supported direct molecular mechanism within the tested system; and organism-level phenotyping supported consequences at the measured biological scale. Functional interpretation of insect small RNAs particularly illustrates why these evidence classes should remain distinct [1]. Transcript-protein modeling likewise demonstrates that apparent molecular discordance should be tested against simpler kinetic explanations before invoking additional regulatory layers [4].

Network construction followed the same boundary logic. An edge between regulatory states was considered evidence-supported only when a selected study directly connected the states within a compatible biological system. Methodological work on A-to-I detection was used to define an editing-validation boundary rather than to infer insect biology [15]. Integrated binding, perturbation, and translational analysis was treated as stronger evidence for functional microRNA targeting than sequence prediction alone [21]. Productive-translation measurements were distinguished from ribosome occupancy when defining translational output [28].

Cross-study relationships were not converted into causal edges. Instead, they formed the basis of the proposed architecture only when the relationship was biologically coherent and remained visibly identified as synthesis. No quantitative network weights, certainty scores, universal pathway direction, or predictive-performance estimates

were assigned. The resulting architecture is therefore an evidence-bounded conceptual tool for comparing post-transcriptional states, not an empirically validated predictive model

Conclusion

Post-transcriptional RNA regulation extends insect molecular control beyond measurements of transcript abundance by altering transcript persistence, isoform identity, nucleotide state, small-RNA targeting, chemical modification, and translational realization. These mechanisms can interact, but the available evidence supports a mixture of independent regulation and context-specific cross-layer coupling rather than one universal pathway. The proposed state architecture separates molecular detection from mechanism, organismal consequence, and higher-order ecological or evolutionary interpretation. Its principal value is therefore inferential discipline: it identifies where stronger conclusions require additional evidence and where distinct RNA processes should remain analytically separate. Broader taxonomic coverage, matched multi-layer measurements, functional perturbation, and explicit testing of alternative explanations will be necessary to determine how widely integrated post-transcriptional control operates across insect development and environmental responses.

Acknowledgments: None

Conflict of Interest: None

Financial Support: None

Ethics Statement: None

References

1. Reynolds JA. MicroRNAs in the developmental toolbox - a comparative approach to understanding their role in regulating insect development. Curr Opin Insect Sci. 2024;66:101256. doi:10.1016/j.cois.2024.101256
2. Song J, Zhou S. Post-transcriptional regulation of insect metamorphosis and oogenesis. Cell Mol Life Sci. 2020;77(10):1893-909. doi:10.1007/s00018-019-03361-5
3. Tian S, Monteiro A. A transcriptomic atlas underlying developmental plasticity of seasonal forms of Bicyclus anynana butterflies. Mol Biol Evol. 2022;39(6):msac126. doi:10.1093/molbev/msac126
4. Becker K, Bluhm A, Casas-Vila N, Dinges N, Dejung M, Sayols S, et al. Quantifying post-transcriptional regulation in the development of Drosophila melanogaster. Nat Commun. 2018;9(1):4970. doi:10.1038/s41467-018-07455-9
5. Stewart RK, Nguyen P, Laederach A, Volkan PC, Sawyer JK, Fox DT. Orb2 enables rare-codon-enriched mRNA expression during Drosophila neuron differentiation. Nat Commun. 2024;15(1):5270. doi:10.1038/s41467-024-48344-8
6. Pekovic F, Rammelt C, Kubíková J, Metz J, Jeske M, Wahle E. RNA binding proteins Smaug and Cup induce CCR4-NOT-dependent deadenylation of the nanos mRNA in a reconstituted system. Nucleic Acids Res. 2023;51(8):3950-70. doi:10.1093/nar/gkad159
7. Zhang G, Xu Y, Wang X, Zhu Y, Wang L, Zhang W, et al. Dynamic FMR1 granule phase switch instructed by m6A modification contributes to maternal RNA decay. Nat Commun. 2022;13(1):859. doi:10.1038/s41467-022-28547-7
8. Faucillion ML, Johansson AM, Larsson J. Modulation of RNA stability regulates gene expression in two opposite ways: Through buffering of RNA levels upon global perturbations and by supporting adapted differential expression. Nucleic Acids Res. 2022;50(8):4372-88. doi:10.1093/nar/gkac208
9. Steward RA, de Jong MA, Oostra V, Wheat CW. Alternative splicing in seasonal plasticity and the potential for adaptation to environmental change. Nat Commun. 2022;13(1):755. doi:10.1038/s41467-022-28306-8
10. Netschitailo O, Raub S, Kaftanoglu O, Beye M. Sexual diversification of splicing regulation during embryonic development in honeybees (Apis mellifera), A haplodiploid system. Insect Mol Biol. 2022;31(2):170-6. doi:10.1111/imb.12748

11. Tong C, Zhang K, Rong Z, Mo W, Peng Y, Zheng S, et al. Alternative splicing of POUM2 regulates embryonic cuticular formation and tanning in Bombyx mori. Insect Sci. 2023;30(5):1267-81. doi:10.1111/1744-7917.13164
12. Zhang Z, Bae B, Dredge WH, Miura P. Coordination of alternative splicing and alternative polyadenylation revealed by targeted long read sequencing. Nat Commun. 2023;14(1):5506. doi:10.1038/s41467-023-41207-8
13. Srivastava D, de Toledo M, Manchon L, Tazi J, Juge F. Modulation of Yorkie activity by alternative splicing is required for developmental stability. EMBO J. 2021;40(3):e104895. doi:10.15252/embj.2020104895
14. Duan Y, Ma L, Liu J, Liu X, Song F, Tian L, et al. The first A-to-I RNA editome of hemipteran species Coridius chinensis reveals overrepresented recoding and prevalent intron editing in early-diverging insects. Cell Mol Life Sci. 2024;81(1):136. doi:10.1007/s00018-024-05175-6
15. Wei Q, Han S, Yuan K, He Z, Chen Y, Xi X, et al. Transcriptome-wide profiling of A-to-I RNA editing by Slic-seq. Nucleic Acids Res. 2023;51(16):e87. doi:10.1093/nar/gkad604
16. Yablonovitch AL, Fu J, Li K, Mahato S, Kang L, Rashkovetsky E, et al. Regulation of gene expression and RNA editing in Drosophila adapting to divergent microclimates. Nat Commun. 2017;8(1):1570. doi:10.1038/s41467-017-01658-2
17. Deng P, Khan A, Jacobson D, Sambrani N, McGurk L, Li X, et al. Adar RNA editing-dependent and -independent effects are required for brain and innate immune functions in Drosophila. Nat Commun. 2020;11(1):1580. doi:10.1038/s41467-020-15435-1
18. Örkenby L, Skog S, Ekman H, Gozzo A, Kugelberg U, Ramesh R, et al. Stress-sensitive dynamics of miRNAs and Elba1 in Drosophila embryogenesis. Mol Syst Biol. 2023;19(5):e11148. doi:10.15252/msb.202211148
19. Inui T, Sezutsu H, Daimon T. MicroRNA let-7 is required for hormonal regulation of metamorphosis in the silkworm, Bombyx mori. Insect Biochem Mol Biol. 2022;145:103784. doi:10.1016/j.ibmb.2022.103784
20. Liu Z, Ling L, Xu J, Zeng B, Huang Y, Shang P, et al. MicroRNA-14 regulates larval development time in Bombyx mori. Insect Biochem Mol Biol. 2018;93:57-65. doi:10.1016/j.ibmb.2017.12.009
21. Wessels HH, Lebedeva S, Hirsekorn A, Wurmus R, Akalin A, Mukherjee N, et al. Global identification of functional microRNA-mRNA interactions in Drosophila. Nat Commun. 2019;10(1):1626. doi:10.1038/s41467-019-09586-z
22. Guo Z, Bai Y, Zhang X, Guo L, Zhu L, Sun D, et al. RNA m6A methylation suppresses insect juvenile hormone degradation to minimize fitness costs in response to a pathogenic attack. Adv Sci (Weinh). 2024;11(6):e2307650. doi:10.1002/advs.202307650
23. Liu J, Huang T, Chen W, Ding C, Zhao T, Zhao X, et al. Developmental mRNA m5C landscape and regulatory innovations of massive m5C modification of maternal mRNAs in animals. Nat Commun. 2022;13(1):2484. doi:10.1038/s41467-022-30210-0
24. Liu SQ, Jia SZ, Tian H, Li YH, Hu KW, Tao JG, et al. Evolution of m6A-related genes in insects and the function of METTL3 in silkworm embryonic development. Insect Mol Biol. 2023;32(3):316-27. doi:10.1111/imb.12832
25. Wang Y, Zhang L, Ren H, Ma L, Guo J, Mao D, et al. Role of Hakai in m6A modification pathway in Drosophila. Nat Commun. 2021;12(1):2159. doi:10.1038/s41467-021-22424-5
26. Zhang T, Xue Y, Su S, Altouma V, Ho K, Martindale JL, et al. RNA-binding protein Nocte regulates Drosophila development by promoting translation reinitiation on mRNAs with long upstream open reading frames. Nucleic Acids Res. 2024;52(2):885-905. doi:10.1093/nar/gkad1122
27. Villa S, Dwivedi P, Stahl A, Hinkle T, Rose CM, Kirkpatrick DS, et al. OTUD6 deubiquitination of RPS7/eS7 on the free 40 S ribosome regulates global protein translation and stress. Nat Commun. 2024;15(1):6873. doi:10.1038/s41467-024-51284-y
28. Patraquim P, Mumtaz MAS, Pueyo JI, Aspden JL, Couso JP. Developmental regulation of canonical and small ORF translation from mRNAs. Genome Biol. 2020;21(1):128. doi:10.1186/s13059-020-02011-5