



Eurasia Specialized Veterinary Publication

Entomological Research Letters

ISSN: 3108-639X

2022, Volume 2, Issue 1, Page No: 12-18

Copyright CC BY-NC-SA 4.0

Available online at: www.esvpub.com/

Investigating *Ocimum Basilicum* Extract Effectiveness Against Mosquito Larvae

Rahul Ramirez¹, Omar P. Zhang^{1*}, Anna Santos¹

¹Department of Insect Science, School of Biological Sciences, Swedish University of Agricultural Sciences, Uppsala, Sweden.

*E-mail ✉ ozhang@hotmail.com

ABSTRACT

Mosquito control is a key method for managing diseases transmitted by these insects, and targeting their early life stages offers several advantages, such as ease of application and greater susceptibility to environmental and chemical factors. This study aimed to investigate the chemical constituents and the larvicidal effects of ethanolic extracts from the leaves and flowers of *Ocimum basilicum* on mosquito larvae. Fresh *O. basilicum* was harvested, dried for one week in the shade, and then extracted using ethanol. They were placed in containers and subjected to bioassays according to WHO guidelines. Mortality rates were recorded after a 24-hour recovery period, and dead larvae were examined for any morphological changes. Probit analysis was performed using SPSS version 19. Phytochemical analysis of *O. basilicum* leaves revealed the presence of flavonoids, tannins, glycosides, and steroids, while saponins, alkaloids, and terpenoids were absent. For the flowers, tannins, steroids, terpenoids, and flavonoids were found, but saponins, glycosides, and alkaloids were not present. The LC₅₀ values for the leaf and flower extracts were found to be 17.78 ppm, 16.98 ppm, 15.48 ppm, and 15.84 ppm, with LC₉₅ values at 56.23 ppm, 64.56 ppm, 66.06 ppm, and 50.11 ppm, respectively, for *Anopheles arabiensis* and *Culex quinquefasciatus*. Larvae treated with ethanolic extracts showed visible morphological changes, including discoloration and deformities in the digestive tract.

Keywords: Ethanol extract, Larvicidal, *Anopheles arabiensis*, *Ocimum basilicum*, *Culex quinquefasciatus*, Phytochemicals

Received: 09 March 2022

Revised: 28 April 2022

Accepted: 30 April 2022

How to Cite This Article: Ramirez R, Zhang OP, Santos A. Investigating *Ocimum Basilicum* Extract Effectiveness Against Mosquito Larvae. Entomol Res Lett. 2022;2(1):12-8.

<https://doi.org/10.51847/Lpt5VNYCTdSV>

Introduction

Mosquitoes, such as *Anopheles*, *Culex pipiens*, and *Aedes*, that belong to the Diptera: Culicidae family, are vectors for a variety of serious diseases. The use of chemical insecticides, while effective, is costly and poses significant environmental risks. Furthermore, the growing resistance of many mosquito species to these chemicals has become a major concern [1-3]. In recent years, there has been a growing focus on finding new, naturally occurring compounds from plants to replace chemical insecticides. Plants have proven to be valuable sources of novel natural insecticides. Many plant-based mosquito repellents and insecticides have shown promising results as effective inhibitors of mosquito populations [4-9].

Ocimum basilicum L., commonly known as basil and belonging to the Lamiaceae family, is widely utilized in traditional medicine worldwide. *Aedes aegypti* has shown a high sensitivity to extracts from *O. basilicum* leaves, especially at concentrations ranging from 0.3% to 1.5%. When ethanol is used to extract basil leaf powder, its toxicity to insects increases, making it a recommended solution for controlling pests like the American cockroach [10-12]. Additionally, basil essential oils are particularly effective against the 3rd instar larvae of *A. aegypti*, yielding favorable LC₅₀ and LC₉₀ values in laboratory tests. These oils have also demonstrated 100% mortality

rates while tested against mites. Furthermore, basil leaves have proven effective in controlling agricultural pests in rice, with high mortality rates observed [9-11]. This study aims to analyze the chemical components and evaluate the larvicidal properties of ethanolic extracts from the leaves and flowers of *O. basilicum*, contributing to the exploration of natural alternatives for mosquito control.

Materials and Methods

Plant collection

Fresh *O. basilicum* plants were collected from the garden of the College of Science at Imam Mohammed Ibn Saud Islamic University. The leaves were identified and processed by a faculty member. After collection, the plant parts were air-dried in a shaded area for one week. Once dried, they were ground into a fine powder and stored in sealed plastic containers until needed for further analysis and experiments.

Phytochemical analysis of extracts

To analyze the phytochemical composition, selected parts of *O. basilicum* (leaves and flowers) were subjected to preliminary screening using methods described by Mohamed Nour [13]. This screening was conducted to detect the presence of key chemical compounds, including alkaloids, tannins, saponins, flavonoids, steroids, glycosides, and terpenoids.

Preparation of ethanolic extracts

Ethanolic extracts were prepared by soaking the powdered plant parts in ethanol, following the method established by Hamid *et al.* [14]. The ethanol was allowed to extract the compounds for further analysis.

Mosquito larvae

Mosquito larvae, specifically *A. arabiensis*, and *C. quinquefasciatus*, were collected and cultured under controlled laboratory conditions at the Biology Department, Faculty of Science. The larvae were placed in plastic containers and provided with the required nutrients for growth and development.

Larval bioassay

The larvicidal properties of the ethanolic extracts were tested according to the standard procedure approved by the World Health Organization (WHO) [15]. A set of 20 third- and early fourth-instar larvae from *A. arabiensis* and *C. quinquefasciatus* was placed in 250 ml of tap water in individual plastic cups. The extracts were tested at varying concentrations, and each concentration was replicated three times. Control groups, that did not receive any extract, were also included. Mortality was observed after 24 hours, and the dead larvae were analyzed for morphological changes under a microscope.

Statistical analysis

The data collected from the bioassay were calculated using SPSS version 19. Probit analysis was used to determine the LC50 and LC90 values for the larvae of *A. arabiensis* and *C. quinquefasciatus*. Additional statistical parameters, such as the regression coefficient (R^2), slope, and x-coefficient, were calculated to assess the efficacy of the extracts.

Results and Discussion

Phytochemical Composition of *O. basilicum* Leaves and Flowers

The phytochemical analysis of *O. basilicum* leaves and flowers revealed varying amounts of different compounds (Table 1).

Table 1. Phytochemical composition of *O. basilicum* leaves and flowers

Plant Part	Alkaloids	Saponins	Tannins	Flavonoids	Glycosides	Steroids	Terpenoids
<i>O. basilicum</i> Leaves	-	-	+	+	+	+	-
<i>O. basilicum</i> Flowers	-	+	-	+	++	-	+

- “-” indicates the absence of the compound.
- “+” indicates the presence of the compound.

• “++” indicates the compound is present in relatively higher amounts.

Effect of ethanolic extracts of basil leaves and flowers on mosquito larvae (24 Hours)

The findings of the present study demonstrated that the ethanolic extract from basil leaves, when tested at varying concentrations, resulted in a lethal concentration (LC50) of 17.78 ppm for *Anopheles arabiensis* and 16.98 ppm for *Culex quinquefasciatus* larvae. On the other hand, the ethanolic extract from basil flowers showed an LC50 of 15.48 ppm for *Anopheles arabiensis* and 15.84 ppm for *Culex quinquefasciatus* larvae (Tables 2 and 3; Figures 1 and 2).

The LC50 data revealed that the flower extract exhibited a stronger effect on *Anopheles* larvae than on *Culex* larvae. Moreover, the flower extract demonstrated greater efficacy and biological activity than the leaf extract.

Table 2. The action of ethanol extract of leaves on mosquito larvae (24 hrs)

Conc. (ppm)	Log- Conc	<i>A. arabiensis</i>		<i>C. quinquefasciatus</i>	
		Mortality (%)	Probit	Mortality (%)	Probit
59.3	1.773	97	6.88	92	6.41
47.44	1.676	82	5.92	87	6.13
35.58	1.551	75	5.07	80	5.84
23.72	1.375	57	5.18	52	5.05
11.86	1.074	47	4.92	32	4.53
R2		0.64		0.97	
slope		2.36		2.79	
x-coefficient		2.07		1.43	
LC50		17.78 ppm		16.98 ppm	
LC95		56.23 ppm		64.56 ppm	




Figure 1. Log- Probit curve of action of ethanol extract of leaves on mosquito larvae (24hrs)

Table 3. The action of ethanol extract of flower on mosquito larvae (24hrs)

Conc. (ppm)	Log- Conc	<i>A. arabiensis</i>		<i>C. quinquefasciatus</i>	
		Mortality (%)	Probit	Mortality (%)	Probit
58.56	1.76	90	6.28	97	6.88
46.86	1.67	85	6.04	87	6.13
35.15	1.54	72	5.58	77	5.74
23.43	1.36	57	5.18	62	5.31
5.86	0.76	35	4.61	30	4.48

R2	0.912	0.88
slope	1.60	2.12
x-coefficient	3.25	2.69
LC ₅₀	15.48 ppm	15.84 ppm
LC ₉₅	66.06 ppm	50.11 ppm




Figure 2. Log- Probit curve of action of ethanol extract of flower on mosquito larvae (24hrs)

Damage to mosquito larvae caused by basil plant extracts

Mosquito larvae exposed to basil plant extracts exhibited several deformities. Notable changes included discoloration (bright coloration), detachment of the alimentary canal, and swelling of the alimentary canal, which was not properly connected to the head (**Figure 3**).




Figure 3. The damage to mosquito larvae caused by basil plant extracts. a) Control larva, b) Bright color larva, c) Larvae swollen alimentary canal that was not attached to the head, d) Larva disconnected the alimentary canal

Mosquitoes, recognized as key vectors for transmitting diseases, present a significant challenge in both public health and entomological research [16]. Recently, there has been growing interest in the use of plant-based products as alternatives for pest control. Unlike synthetic insecticides, plant extracts offer a rich source of bioactive compounds that are generally less toxic to mammals and biodegradable [17, 18]. This shift is crucial due to concerns over the increasing resistance of mosquitoes to synthetic chemicals and the environmental and food safety risks posed by traditional insecticides [19].

Ocimum basilicum, known for its diverse phytochemical content, is rich in compounds such as terpenoids, flavonoids, alkaloids, tannins, and glycosides, which contribute to its pharmacological benefits [20]. In this study, the phytochemical analysis of *O. basilicum* leaves revealed the presence of glycosides, tannins, steroids, and flavonoids, with alkaloids, saponins, and terpenoids absent. In contrast, the flowers contained tannins, steroids, terpenoids, and flavonoids, but lacked alkaloids, glycosides, and saponins.

The study found that the phytochemicals in basil have significant larvicidal effects on mosquito larvae. The findings are in line with the research of Azhari *et al.* [9], which demonstrated the larvicidal and repellent properties of *O. basilicum* against the dengue vector (*A. aegypti*). Basil has also been shown to have insecticidal properties against *Culex quinquefasciatus* larvae under laboratory conditions, with aqueous plant extracts exhibiting effective LC50 values of 5.32% [21]. These results further support the potential of basil as an effective mosquito control agent when used in water extracts and other solvents.

In another study, hexane extracts of basil leaves were tested against *Anopheles arabiensis* larvae, demonstrating excellent repellent activity against adult mosquitoes for up to two hours [22]. Although the current study focused on larval stages, the efficacy of ethanol extracts corroborates previous findings. Ethanol extracts of *O. basilicum* exhibited significant repellency and larvicidal activity against *A. aegypti* and *C. quinquefasciatus*, supporting the consistency of these results with prior studies [23].

Furthermore, basil's essential oils, when used as larvicides, show promising effects against *Anopheles arabiensis* larvae, with an LC50 value of 58 mg/l and LC90 of 143 mg/l, contributing to malaria control efforts [24]. These findings align with the current study's results and emphasize the potential of plant-based bio-agents in mosquito control programs. Many studies have suggested that essential oils of *O. basilicum* could serve as a natural repellent [25-27], with *Anopheles* larvae proving more susceptible than *Culex* species [28]. Basil leaf essential oils have also been recommended as an efficient repellent and a moderate larvicide against *A. aegypti* [29, 30].

The current study also highlighted the morphological changes induced by basil extracts in mosquito larvae, including discoloration, swelling, and detachment of the alimentary canal. These deformities are consistent with observations in other studies, where plants like *L. camara* and *A. indica* caused similar morphological changes in mosquito larvae [14, 31]. These findings confirm the effectiveness of *O. basilicum* as a potential bio-pesticide for mosquito control.

Conclusion

The study found that the ethanol extract of *O. basilicum* demonstrated greater effectiveness against *Anopheles arabiensis* larvae compared to *Culex quinquefasciatus* larvae, with the flower extract showing higher mortality rates than the leaf extract. Based on these findings, the study recommends *O. basilicum* ethanol extract as an effective, eco-friendly natural larvicide.

Acknowledgments: None

Conflict of Interest: None

Financial Support: None

Ethics Statement: None

References

1. El Ouali Lalami A, El-Akhal F, Ez Zoubi Y, Taghzouti K. Study of phytochemical screening and larvicidal efficacy of ethanol extract of *Salvia officinalis* (Lamiaceae) from north center of Morocco against *Culex*

- piapiens (Diptera: Culicidae) vector of serious human diseases. *Int J Pharmacog Phytochem Res.* 2016;8(10):1663-8. Available from: www.ijpr.com
2. Krishnappa K, Pandiyan J, Elumalai K, Baranitharan M, Jayakumar S, Gokulakrishnan J. GC–MS analysis and mosquitocidal properties of *Loranthus pentandrus* Linn. (Loranthaceae) against human vector mosquitoes (Diptera: Culicidae). *Acad J Med Plants.* 2019;7(12):261-8. doi:10.15413/ajmp.2019.0154
 3. Lupi E, Hatz C, Schlagenhauf P. The efficacy of repellents against *Aedes*, *Anopheles*, *Culex* and *Aedes* spp. a literature review. *Trav Med Infect Dis.* 2013;11(6):374-411.
 4. Asadollahi A, Khoobdel M, Zahraei-Ramazani A, Azarmi S, Mosawi SH. Effectiveness of plant-based repellents against different *Anopheles* species: a systematic review. *Malar J.* 2019;18(1):1-20. doi:10.1186/s12936-019-3064-8
 5. Sukumar K, Perich MJ, Boobar LR. Botanical derivatives in mosquito control: a review. *J Am Mosq Control Assoc.* 1991;7(2):210-37.
 6. Miele M, Dondero R, Ciarallo G, Mazzei M. Methyleugenol in *Ocimum basilicum* L. Cv. genovese gigante. *J Agric Food Chem.* 2001;49(1):517-21.
 7. Husna I, Setyaningrum E, Handayani TT, Kurnia Y, Palupi EK, Umam R, et al. Utilization of basil leaf extract as anti-mosquito repellent: a case study of total mosquito mortality (*Aedes aegypti* 3rd Instar). *J Phys: Conf Ser.* 2020;1467(1):012014.
 8. Ukoroije BR, Island W, Island W, Island W. The efficacy of *Ocimum gratissimum* leaf powder and ethanol extract on adult *Periplaneta Americana* under laboratory condition. *Open Access Libr J.* 2018;5(04):1. doi:10.4236/oalib.1104455
 9. Azhari H, Abdurahman H, Mashitah M, DO SJ. Bioactive compounds from Basil (*Ocimum basilicum*) essential oils with larvicidal activity against *Aedes aegypti* larvae. 3rd Int. Conf Bio, Env Chem, Singapore. 2012;46:21-4.
 10. Amer A, Mehlhorn H. Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitol Res.* 2006;99(4):466-72.
 11. Perumalsamy H, Kim JY, Kim JR, Hwang KN, Ahn YJ. Toxicity of basil oil constituents and related compounds and the efficacy of spray formulations to *Dermatophagoides farinae* (Acari: Pyroglyphidae). *J Med Entomol.* 2014;51(3):650-7.
 12. López MD, Jordán MJ, Pascual-Villalobos MJ. Toxic compounds in essential oils of coriander, caraway and basil active against stored rice pests. *J Stored Prod Res.* 2008;44(3):273-8.
 13. Mohamed Nour AA. Chemical component of the essential oil of basil (*Ocimum basilicum* L.) active against *Salmonella typhi* [Doctoral dissertation, M. Sc. Thesis]. Sudan: University of Gezira; 2009.
 14. Hamid NS, Kehail MA, Ibrahim NA, Abdel-Rahman EH. Larvicidal activity of ethanol extracts of *Azadirachta indica* (Neem) against *Anopheles Arabiensis* and *Culex quinquefasciatus* larvae, Gezira state, Sudan. *Int J Entomol Res.* 2021;6(1):138-41.
 15. World Health Organization. Pesticides and their application: for the control of vectors and pests of public health importance. Geneva: World Health Organization; 2006. Report No.: WHO/CDS/WHOPES/GCDPP/2006.1.
 16. Yousif HY. Susceptibility of *Anopheles arabiensis* Patton and *Culex quinquefasciatus* say larvae to *Ixora* (*Ixora coccinea* L.) leaves and flowers extracts [dissertation on the internet]. Sudan: university of Gezira; 2013. [cited 2013 May 16]. Available from: <http://repo.uofg.edu.sd/handle/123456789/2165>
 17. Aouati A, Roubhi AH, Boudjahem I, Berchi S. Study of toxicological activity of the essential oil of *Ocimum basilicum* L. against Culicidae larvae in Algeria. *AIP Conf Proc.* 2019;2190(1):020073. doi:10.1063/1.5138559
 18. Madhumitha G, Rajakumar G, Roopan SM, Rahuman AA, Priya KM, Saral AM, et al. Acaricidal, insecticidal, and larvicidal efficacy of fruit peel aqueous extract of *Annona squamosa* and its compounds against blood-feeding parasites. *Parasitol Res.* 2012;111(5):2189-99. doi:10.1007/s00436-011-2671-2
 19. Shaalan EA, Canyon D, Younes MW, Abdel-Wahab H, Mansour AH. A review of botanical phytochemicals with mosquitocidal potential. *Environ Int.* 2005;31(8):1149-66. doi:10.1016/j.envint.2005.03.003
 20. Al-Snafi AE. Chemical constituents and pharmacological effects of *Ocimum basilicum*-a review. *Int J Pharm Res.* 2021;13(2):2997-3013.

21. Iqbal J, Ishtiaq F, Alqarni AS, Owayss AA. Evaluation of larvicidal efficacy of indigenous plant extracts against *Culex quinquefasciatus* (Say) under laboratory conditions. *Turk J Agric For.* 2018;42(3):207-15. doi:10.3906/tar-1711-69
22. Elsiddig FEI, Khei SEM. Effect of Neem, Cafure, and Rehan organic extracts compared to other chemical and biological standard insecticides on *Anopheles arabiensis* Patton mosquito. [dissertation on the internet]. Agricultural studies college of agricultural studies: Sudan University of science and technology; 2007. [cited 2007 April 1]. Available from: <http://repository.sustech.edu/handle/123456789/3504>
23. Phasomkusolsil S, Soonwera M. Comparative mosquito repellency of essential oils against *Aedes aegypti* (Linn.), *Anopheles dirus* (Peyton and Harrison) and *Culex quinquefasciatus* (Say). *Asian Pac J Trop Biomed.* 2011;1(1):S113-8. doi:10.1016/S2221-1691(11)60136-6
24. Ali HE. Effect of basil (*Ocimum basilicum* L.) leave powder and ethanolic extract on the 3rd larval instar of *Anopheles arabiensis* (Patton,1905) Culicidae: Diptera. [dissertation on the internet]. Sudan: university of Gezira; 2013. [cited 2013 January 31]. Available from: <http://repo.uofg.edu.sd/handle/123456789/2171>
25. Panneerselvam C, Murugan K, Kovendan K, Kumar PM, Subramaniam J. Mosquito larvicidal and pupicidal activity of *Euphorbia hirta* Linn. (Family: Euphorbiaceae) and *Bacillus sphaericus* against *Anopheles stephensi* Liston.(Diptera: Culicidae). *Asian Pac J Trop Med.* 2013;6(2):102-9. doi:10.1016/S1995-7645(13)60003-6
26. Ileke KD, Adesina JM. Toxicity of *Ocimum basilicum* and *Ocimum gratissimum* extracts against main malaria vector, *Anopheles gambiae* (Diptera: Culicidae) in Nigeria. *J Arthropod Borne Dis.* 2019;13(4):362.
27. Adam AA, Ahmed SA, Mohamed TA, Azrag RA, Mustfa SE, Hamdi AA. Evaluation of repellent activities of the essential oil of *Ocimum basilicum* against anopheles mosquito and formulation of mosquitoes repellent cream. *Biomed Res Clin Prac.* 2019;4(2):2-5. doi:10.15761/BRCP.1000184
28. Abdalla AI, Kehail MA, Abdelrahim YM, Ibrahim NA. Phytochemical screening of *Calotropis procera* ait flower parts and their larvicidal potentialities against anopheles and *Culex* larvae, Gezira state, Sudan. *Int J Biol Res.* 2017;2(2):88-92.
29. Govindarajan M, Rajeswary M. Ovicidal and adulticidal potential of leaf and seed extract of *Albizia lebbek* (L.) Benth. (Family: Fabaceae) against *Culex quinquefasciatus*, *Aedes aegypti*, and *Anopheles stephensi* (Diptera: Culicidae). *Parasitol Res.* 2015;114(5):1949-61. doi:10.1007/s00436-015-4384-4
30. Kumar S, Warikoo R, Mishra M, Samal RR. Impact of *Ocimum basilicum* leaf essential oil on the survival and behaviour of an Indian strain of dengue vector, *Aedes aegypti* (L.). *Vector Biol J.* 2017;2:2. doi:10.4172/2473-4810.1000122
31. Alghamdi AA, Basher NS. Efficacy of leaves and flowers ethanol extracts of the invasive species *lantana camara* Linn as a mosquito larvicidal. *Int J Mosq Res.* 2020;7(5):43-7.