



Eurasia Specialized Veterinary Publication

Entomological Research Letters

ISSN:3108-639X

2021, Volume 1, Issue 2, Page No: 15-20

Copyright CC BY-NC-SA 4.0

Available online at: www.esvpub.com/

Beetle Species Associated with Nesting Birds of the North Caucasus

James Rossi^{1*}, Emily Nakamura¹

¹Department of Insect Ecology, School of Biological Sciences, University of Bologna, Bologna, Italy.

*E-mail ✉ jrossi@outlook.com

ABSTRACT

This paper provides a faunistic inventory of Coleoptera species from 13 families, including 40 species collected from the nests of 24 bird species in the North Caucasus. Compared to our previous data, the list of beetle species has been expanded by 2 more species. Around 20 species were identified as accidental visitors to the bird nests, but these are not discussed further in this paper. The remaining species show intricate biocenotic interactions with their host nests. The process of adaptation of species to the nest environment, a form of microbiocenosis, represents a significant evolutionary development that requires further investigation. Geographically, the North Caucasus serves as a refugium for birds, with many species remaining year-round. As a result, the biodiversity of nidicolous Coleoptera is notably higher in this region compared to more northern areas. The data presented here has contributed to a database of beetle species inhabiting bird nests, which will facilitate future studies on the seasonal dynamics of populations and their associations with specific hosts. This research is particularly relevant in light of recent findings on climate change and human impact on natural ecosystems. As bird species decline and the prevalence of synanthropic species increases, it is possible to predict changes in the populations of nidicolous species, some of which have significant sanitary and epidemiological importance.

Keywords: Coleoptera, Birds nest, Nidicolous, Beetles

Received: 08 September 2021

Revised: 11 November 2021

Accepted: 16 November 2021

How to Cite This Article: Rossi J, Nakamura E. Beetle Species Associated with Nesting Birds of the North Caucasus. Entomol Res Lett. 2021;1(2):15-20.
<https://doi.org/10.51847/MPAJchaG2yj>

Introduction

Beetles were among the first non-parasitic arthropods to be studied concerning nidicolous animal fauna [1-10]. As early as the previous century, beetles were identified as inhabitants of bird nests [1, 2, 4-8, 11-13]. Initial efforts to examine beetles in bird nests and explore the biological connections between the nest owners and their nest-dwelling residents were made in the works of various researchers [1-7, 14-19]. Our contributions to this field have been published in several articles [20-25].

Currently, over 200 beetle species have been recorded with known infestations of helminth larvae [3-7, 14-16, 26], increasing the focus not only on identifying beetle species but also on studying their role in the nest microbiocenosis. Research has shown that Dermestidae species can serve as vectors for pathogen transmission [8, 24, 25]. Gamasid mites often found phoretically attached to necrobiont beetles, have been identified in our studies, including *Poecilochirus necrophori* and *Parasitus* spp., which are capable of carrying rickettsioses [24, 25].

A promising avenue for further research could involve examining the role of birds as vectors for the spread of non-native beetle species, such as Dermestidae, that are associated with bird nests. This could include comparing the primary habitats of these beetles, observed as phoronts, with bird migration patterns. This article provides a faunistic inventory of Coleoptera species from 13 families, comprising 40 species collected from the nests of 24 bird species in the North Caucasus.

Materials and Methods

The material was collected from various regions in the North Caucasus, including the Stavropol Territory (Novoaleksandrovsk, Krasnogvardeisky, Izobilnensky, Shpakovsky, Kochubeevsky, Andropovsky, Grachevsky, Mineralovodsky, Predgorny, Levokumsky, Ipatovsky districts), Krasnodar Territory (Novopavlovka village), Rostov Region (Rostov-on-Don, Rasypnoe village, Zhukovskoe village), North Ossetia (Mozdok town), and Kalmykia (Khurulyun estuary, Tsaryk lake, Solenoe village, Lake Manych). Stationary research was conducted in the Stavropol Upland region between 2000 and July 2020. This study presents data on beetle species from 13 families, which form the core of the nidicolous beetle fauna in the study area. Other coleopteran species found in the nests are not included in this work but are considered nidicolophilic factors in the biology of these species. To better understand which bird species are most frequently visited by Coleoptera, we have reviewed various ornithological studies [27-37].

Results and Discussion

It should be emphasized that environmental pollution contributes to a decline in the population of nesting birds, which in turn negatively impacts the beetle species that are nidicolous [11, 12, 19, 26-29, 38-40]. Beetles were collected from bird nests (**Table 1**), with a total of 390 nests examined. In all, 2,711 beetles, three hundred fifty larvae, and 42 pupae were gathered. The distribution of species across host nests is presented in **Table 1**. In certain instances, the presence of insects in the nests is difficult to explain and is likely due to random factors. When classifying the identified beetles by their feeding habits, it was found that 24 species are polyphages, 20 are necrophages, 20 are keratophages, 16 are coprophages, 20 are detritophages, and 8 are saproxylophages. The trophic preferences of preimaginal stages may vary, with cannibalism being typical in species from the genera *Thanatophilus* and *Dermestes* [19, 25].

Table 1. The confinement of beetles to bird nests (North Caucasus)

| List of beetle species | Host species in a systematic manner | | | | | | | | | | | | | | | | | | | | | | | | |
|-----------------------------------|-------------------------------------|---------------------------|------------------------|-------------------------|---------------------------|-----------------------|----------------------|----------------------|------------------------------|------------------|--------------------|-----------------------|------------------------------|------------------------|--------------------------|------------------------|------------------|--------------------------|------------------|------------------------|--------------------------|--------------------|--------------------------|----------------------|---|
| | <i>Anser fabalis</i> | <i>Anas platyrhynchos</i> | <i>Pernis apivorus</i> | <i>Circus macrourus</i> | <i>Accipiter gentilis</i> | <i>Falco subbuteo</i> | <i>Perdix perdix</i> | <i>Columba livia</i> | <i>Streptopelia decaocto</i> | <i>S. turtur</i> | <i>Otus scopus</i> | <i>Merops apister</i> | <i>Pyonoprogne rupestris</i> | <i>Hirundo rustica</i> | <i>Galerida cristata</i> | <i>Lanius collurio</i> | <i>Pica pica</i> | <i>Corvus frugilegus</i> | <i>C. cornix</i> | <i>Sylvia communis</i> | <i>Passer domesticus</i> | <i>P. montanus</i> | <i>Emberiza calandra</i> | <i>E. citrinella</i> | |
| Silphidae | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Nicrophorus vespillo</i> L. | | | + | | + | | | | | + | | | | | | | + | | | | | | | | |
| <i>N. fossor</i> Er. | | | + | | | | | | | | | | | | | | | | | | | | | | |
| <i>N. vespilloides</i> Hbst. | | | | | | + | | | | + | | | | | | | + | | | | | | | | |
| <i>Silpha obscura</i> L. | | | | | + | | | + | | | | | | | | | | | | | | | | | |
| <i>Silpha obscura</i> (larva) | + | | | | | | | | + | | | | | + | + | + | | + | | | | + | | + | |
| <i>S. carinata</i> Hbst. | | + | | | | + | | | | | | | | | | | + | | | | | | | | |
| <i>S. carinata</i> (larva) | + | | | | | | | | + | | | | | | | | + | | | + | | | + | | |
| <i>Tanatophilus sinuatus</i> F. | | | + | | | + | | | | + | | | + | | | | | | | | | | | | |
| <i>T. sinuatus</i> (larva) | | | | + | | + | + | + | + | | | | | | | | | | + | | | | | | |
| <i>T. terminatus</i> Humm. | | | | | + | | + | | | | | | | | | | | | | | | | | | |
| Dermestidae | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Dermestes frischii</i> (larva) | | + | | | | + | + | + | | + | + | + | + | + | + | + | + | + | + | + | | | + | + | + |
| <i>D. bicolor</i> (larva) | | | | | | + | | | | + | + | | + | | + | + | + | + | + | + | + | | | | |
| <i>D. bicolor</i> (pup.) | | | | | | + | | | | + | | | | | + | + | + | + | + | + | + | | | | |
| <i>D. bicolor</i> F. | | | | | | + | | | | + | | + | | | + | + | + | + | + | + | | | | | |
| <i>D. coronatus</i> Stev. | | | | + | | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | | | + | | |

These beetles, both in their adult and larval stages, may enter the chicks' ear and eye openings, leading to their death. The remains of dead chicks and abandoned eggs attract necrophagous beetles like *Nicrophorus* to the nests. While these beetles feed on the carcasses, they do not reproduce in the nests, which has been observed in the nests of large birds of prey (which were not part of our study). Detritivores and saproxylophages feed on the leftover food in the nest, while coprophages consume droppings and bird pellets, and keratophages target feathers, down, and feather sheaths.

As adults, beetles from *Attagenus* and *Anthrenus* genera require additional nourishment from flowering plants. Our nest inspections displayed that the species and number of beetles are primarily influenced by the nest's structure and the types of food available. Dermestidae and Trogidae favor enclosed nests, where larvae are better protected from external factors, with more stable temperatures, less temperature fluctuation, consistent humidity, and improved conditions for winter diapause. In contrast, Silphidae and Catopidae are more common in open nests near bodies of water. Scarabaeidae can be found in various nest types. The material used to construct the nest also plays a key role. Dermestidae do not inhabit nests made from plant matter, except in birds of prey nests. Families such as Ptinidae, Byrrhidae, and Hydrophilidae are part of the saproxyl population, relying on plant-based protein, while species from the Tenebrionidae family can consume animal-based protein.

Nidicolous insects are inherently linked to their host's life cycle, though this relationship is more tightly integrated in species of *Dermestes* than in keratophages (such as *Attagenus*, *Anthrenus*, and *Trox*) and other groups. This stronger bond is due to *Dermestes* species primarily feeding on food brought to the chicks, which is available only while the chicks are in the nest. As a result, the developmental cycle of *Dermestes* beetles coincides with that of the host nest. This is why *Dermestes* species are typically univoltine; their larvae develop while chicks are being fed, and the newly emerged beetles only begin reproducing in the spring of the following year.

In contrast, keratophages and other groups are less reliant on the host's life cycle. Keratophages feed on wool and feathers, which remain in the nest for extended periods, especially in steppe and semi-desert regions. These beetles typically enter the nest soon after its construction and continue to reproduce throughout the warm season. The feeding on keratin allows for the possibility of larvae developing over two years. Another noteworthy group is the Nitidulidae family, which feeds on the food brought to the nest but also actively searches for additional food sources beyond the nest.

At first, it might seem unusual that larvae from different nidicolous species coexist in such confined environments as nests, all developing on the same food. One might expect intense competition among these groups, potentially causing the displacement of one group of beetles. However, a closer examination of the nests, particularly within the Dermestidae and Trogidae families, sheds light on this apparent paradox. It was found that, despite the nests' small size, adult and larval beetles of *Dermestes*, *Anthrenus*, *Attagenus*, and *Trox* species are spatially segregated within the nest. The larvae of *Trox* species inhabit the lower, more compact layers, where their fusiform body shape helps them move. In contrast, *Anthrenus* larvae and adults are more frequently found in the upper, looser layers of the nest substrate, closer to the surface. *Dermestes* beetles occupy the outermost parts of the nest, living along the outer boundaries or within the litter on the nest's walls. *Trox* beetles are found throughout the entire perimeter of the nest. Although beetles from other families might leave the nest temporarily, they always return. Our observations reveal that Catopidae species, such as *Catops picipes* and *C. nigrita*, develop in decaying and spoiled eggs. The adult beetles lay their eggs inside the eggs, and the larvae feed on the egg contents, remaining inside the shell until they pupate. The synanthropic beetle *Dermestes lardarius* is commonly found in urban areas, particularly in the nests of synanthropic birds and in the attics of buildings where the temperature remains above +11°C. It has been observed laying eggs as late as December. Mrochkovsky (1955) reported the significant harm caused by these synanthropic beetles in pigeon and chicken coops in Poland [17]. This species is often found in attics, where it can continue to develop throughout the year. We expect our collection to be expanded further by Latridiidae species, as noted by the writers of the article [37, 41, 42].

Conclusion

The shift to nesting microbiotopes, in our view, involved the development of new ecological niches and the establishment of novel interspecific interactions. This process was characterized by rapid speciation, resulting in the emergence of new ecobiomorphs—nidicolous species that are uniquely adapted to the specific conditions of nests and the food available within them. *Nesting microbiocenoses*, as fundamental systems within the supraorganism framework, provide valuable insight into biocenotic structures at their most basic level.

In the subsequent article, we will explore the mechanisms behind the division of ecological niches among nidicolous organisms, as well as the evolutionary pathways of the nidicolous fauna [33-36].

Acknowledgments: The writers would like to thank the ornithologists who assisted in the collection of material from bird nests: Sabelnikova-Begashvili N.V., Ashibokov U.Kh., Chursina N.V., and Professor Ilyukh M.P. for their assistance in identifying bird species.

Conflict of Interest: None

Financial Support: None

Ethics Statement: This article presents original data collected by the authors through fieldwork and subsequent data processing.

References

1. Vlasov YaP. Nora as a kind of biotope in the vicinity of Ashgabat. *Probl Parasite Fauna of Turkmenistan. Leningrad–Moscow*; 1937. p. 183-9.
2. Vlasov YaP, Shestoporov EL. Beetles from holes in the vicinity of Ashgabat. *Tr Board Stud Produces Force, Ser Turkmen. 1937*;9:269-75.
3. Kirschenblat YaD. On the origin of some species of nidicoles (nest dwellers). *Dokl Akad Nauk SSSR. 1935*;2(3–4):332–7.
4. Negrobov VN. Materials on the parasitization of biohelminth larvae in coprophagous beetles (Coprinae, Coleoptera) in the southeast of the Central Chernozem Belt. In: *Materials of the Scientific Conference of All-Union Society of Helminthologists. 1963*;2:22–3.
5. Polozhentsev PA, Negrobov VP. To the study of insect hosts of biohelminths in the Voronezh region. In: *Works on Helminthology on the 80th Anniversary of Acad. K.I. Skryabin. Moscow*; 1958. p. 274–88.
6. sedenova LU. Necrophilous beetles as intermediate and additional hosts of vertebrate helminths. In: *Collection of Scientific Works: Animal World of Kalmykia, Its Protection and Rational Use. Elista*; 1982. p. 44–50.
7. Joy NH. Coleoptera occurring in the nests of mammals and birds. *Entomol Mon Mag. 1906*;17(42):198-202.
8. Pushkin SV, Belous VN, Alikhadzhev MK, Erzhapova RS, Bagrikova NA. Materials on the fauna of micetobiont and mycetophilic beetles (Coleoptera) Ciscaucasia. *Entomol Appl Sci Lett. 2019*;6(4):1-6.
9. Pushkin SV, Kharchenko LN. Contemporary arahnofauna of Stavropol urban agglomeration. *Entomol Appl Sci Lett. 2017*;4(4):23-6.
10. Sazhnev AS, Matyukhin AV. Cases of unintentional phoresy of beetles (Insecta: Coleoptera) on birds. *Ecosystem Transform. 2019*;2(2):29-33. doi:10.23859/estr-190311
11. Pushkin SV, Kharchenko LN. A variant of the methodology for assessing the state of zoological complexes. *Entomol Appl Sci Lett. 2017*;4(1):23-5.
12. Pushkin S, Tsymbal B, Artemev S, Fesenko G, Fesenko T. Eco-faunistic review of the silphidae family (Coleoptera: Silphidae) of the greater Caucasus. *Entomol Appl Sci Lett. 2020*;7(1):61-6.
13. Pushkin SV. *Insecta: Coleoptera inventory of the Precaucasian and adjacent territories. Moscow (RU): Direct-Media*; 2019. 229 p. doi:10.23681/575393
14. Bickhardt H. Käfer in maulwurfnestern. *Etmomol Blätt. 1916*;12:49-53.
15. Gensicke F. Zur Kenntnis der Nestfauna einiger Muridenarten in der Umgebung von Greifswald. *Wiss Zeitschr Ernst-Moritz-Arndt-Univ Greifswald. 1960*;9(2–3):189–97.
16. Loew F. Bewohner der Schwalbennester. *Verh KK Zool Bot Ges Wien. 1867*;12:237–43.
17. Mroczkowski M. Silphidae. *Klucze do oznaczania owadow Polski. Warszawa. 1955*;19:25-30.
18. Norman J. Coleoptera from old birds nests. *Entomol Month Mag. 1906*;17:39-76.
19. Strouhal H, Beier M. Beitrag zur coleopterenfauna der maulwurfsnester in der nächsten umgebung wiens. *Z Morphol Oekol Tiere. 1928*;12:191-239.
20. Pushkin S, Tsymbal B, Artemev S, Fesenko G, Fesenko T. Eco-faunistic review of the silphidae family (Coleoptera: Silphidae) of the greater caucasus. *Entomol Appl Sci Lett. 2020*;7(1):44-8.

21. Pushkin SV, Tsymbal BM, Rybalova OV. Use of population indicators of beetle (Coleoptera, Silphidae, Dermestidae) in bioindicacion of the environmental status. *Entomol Appl Sci Lett.* 2019;6(4):13-7.
22. Pushkin SV, Tsymbal BM. Ecologic and zoogeographic characteristics of the genus orphilus Er.(Coleoptera: Dermestidae) in Palearctic zone. *Entomol Appl Sci Lett.* 2019;6(3):27-32.
23. Viktorovich PS. Revision of the material of collections of the carpet beetle (Dermestidae). Part 1. Faunistic review. *Entomol Appl Sci Lett.* 2019;6(1):8-21.
24. Pushkin SV. Discoveries of carpet beetles (Coleoptera: Dermestidae) of the south of Russia. *Entomol Appl Sci Lett.* 2017;4(2):29-31. doi:10.24896/easl2017427
25. Pushkin SV. Environmental group necrophilous and necrobionts beetles (Insecta; Coleoptera) of the south of the Russia. *Entomol Appl Sci Lett.* 2015;2(4):1-9.
26. Vojtov VA, Tsymbal BM. Study of tribological characteristics of compatible materials in tribosystems of extruders for manufacturing solid fuel from biomass. *J Friction Wear.* 2018;39(6):500-4.
27. Samplonius JM, Kappers EF, Brands S, Both C. Phenological mismatch and ontogenetic diet shifts interactively affect offspring condition in a passerine. *J Anim Ecol.* 2016;85(5):1255-64.
28. Hamao S, Higuchi M, Jinbo U, Maeto K, Furuki K. Interaction among birds, mosses and insects in bird nests. *Jpn J Ornithol.* 2016;65(1):37-42. 10.3838/jjo.65.37.
29. Biddle LE, Broughton RE, Goodman AM, Deeming DC. Composition of bird nests is a species-specific characteristic. *Avian Biol Res.* 2018;11(2):132-53.
30. Hanmer HJ, Thomas RL, Beswick GJ, Collins BP, Fellowes MD. Use of anthropogenic material affects bird nest arthropod community structure: influence of urbanisation, and consequences for ectoparasites and fledging success. *J Ornithol.* 2017;158(4):1045-59.
31. Radhamany D, Das KS, Azeez PA, Wen L, Sreekala LK. Usage of nest materials by house sparrow (*Passer domesticus*) along an urban to rural gradient in Coimbatore, India. *Trop Life Sci Res.* 2016;27(2):127-34. doi:10.21315/tlsr2016.27.2.10
32. Reynolds SJ, Davies CS, Elwell E, Tasker PJ, Williams A, Sadler JP, et al. Does the urban gradient influence the composition and ectoparasite load of nests of an urban bird species? *Avian Biol Res.* 2016;9(4):224-34. doi:10.3184/175815516X14725499175665
33. Lapshin NV, Matantseva MV, Simonov SA. Choosing a place for a nest and nesting in the willow warbler (*Phylloscopus trochilus* L., 1758) (Sylviidae, Aves) in the taiga zone of north-west Russia. *Povolzhsky Ecol J.* 2019;(3):322-34. doi:10.35885/1684-7318-2019-3-322-334
34. Gullan PJ, Cranston PS. *The insects an outline of entomology.* Wiley- Blackwell; 2017. 632 p. ISBN 978-1-118-84615-5. Available online: <https://trove.nla.gov.au/work/9113436>.
35. Ilyukh MP. Wintering birds of prey and owls of the Ciscaucasia. *Russ Ornithol J.* 2020;29(1931):2516-31.
36. Sazhnev AS, Kondrat'ev EN. Coleoptera (Insecta: Coleoptera) from NOR-swallows beregovushek riparia riparia (Linnaeus, 1758) (Aves: Hirundinidae) field Saratov region. *J Biol.* 2021;2(4):276-81.
37. Vlasov DV, Nikitsky NB, Saluk SV. Fauna of lurking beetles (Coleoptera, Latridiidae) of the Yaroslavl region. *Bull Moscow Soc Nat. Dep Biol.* 2018;123(4):8-18.
38. Vojtov VA, Biekirov AS, Voitov AV, Tsymbal BM. Running-in procedures and performance tests for tribosystems. *J Friction Wear.* 2019;40(5):376-83.
39. Vojtov VA, Kravtsov AG, Tsymbal BM. Evaluation of tribotechnical characteristics for tribosystems in the presence of fullerenes in the lubricant. *J Frict Wear.* 2020;41(6):521-5.
40. Spitsov D, Nekrasova L, Kondratenko L, Pushkin S, Klyuchnikov D. The effect of agricultural practices on the drinking water quality: a case study. *Asian J Water Environ Pollut.* 2020;17(2):73-80. doi:10.3233/AJW200023
41. Pushkin SV. *Necrobiont Coleoptera of the North Caucasus.* Moscow (RU): Direct-Media; 2019. 183 p. doi:10.23681/575396
42. Victorovich PS, Pavlovich IM. Endemism, relicts and invasion species of animals in structure of the biodiversity of the Ciscaucasia. *Entomol Appl Sci Lett.* 2018;5(1):17-20.